

Compass PTN Core Measures

Measure ID	Measure Name	Denominator	Numerator	NQS Domain	Measure Type/ QPP Priority Level	QPP Specialty Measure Set
eMeasure ID: CMS122v5 eMeasure NQF: N/A NQF: 0059 QualityID: 001	Diabetes: Hemoglobin A1c (HbA1c) Poor Control (<9%)	Patients 18-75 years of age with diabetes with a visit during the measurement period.	Patients whose most recent HbA1c level (performed during the measurement period) is >9.0%.	Effective Clinical Care	Intermediate Outcome High Priority	Internal Medicine Preventive Medicine General Practice/Family Medicine
eMeasure ID: CMS165v5 eMeasure NQF: N/A NQF: 0018 QualityID: 236 <i>*This measure is a TCPI Common Measure</i>	Controlling High Blood Pressure	Patients 18-85 years of age by the end of the measurement year who had at least one outpatient encounter with a diagnosis of hypertension (HTN) during the first six months of the measurement year.	The number of patients in the denominator whose most recent BP is adequately controlled during the measurement year. For a patient's BP to be controlled, both the systolic and diastolic BP must be <140/90mmHg (adequate control). To determine if a patient's BP is adequately controlled, the representative BP must be identified.	Effective Clinical Care	Intermediate Outcome High Priority	Internal Medicine Cardiology Obstetrics/ Gynecology Preventive Medicine Thoracic Surgery Vascular Surgery General Practice/Family Medicine
CMS: 126v4 NQF: 0036 PQRS: 311 <i>*not a QPP Metric</i>	Use of Appropriate Medications for Asthma	Patients 5-64 years of age with persistent asthma and a visit during the measurement period.	Patients who were dispensed at least one prescription for a preferred therapy during the measurement period.	Effective Clinical Care	Process Not QPP	
eMeasure ID: CMS144v5 eMeasure NQF: 2908 NQF: 0083 QualityID: 008	Heart Failure (HF): Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction (LVSD)	All patients aged 18 years and older with a diagnosis of heart failure with a current or prior LVEF < 40%.	Patients who were prescribed beta-blocker therapy either within a 12 month period when seen in the outpatient setting OR at each hospital discharge.	Effective Clinical Care	Process	Cardiology Hospitalists General Practice/Family Medicine
eMeasure ID: CMS166v6 eMeasure NQF: N/A NQF: 0052 QualityID: 312 <i>*This measure is a TCPI Common Measure</i>	Use of Imaging Studies for Low Back Pain	Patients 18-50 years of age with a diagnosis of low back pain during an outpatient or emergency department visit.	Patients without an imaging study conducted on the date of the outpatient or emergency department visit or in the 28 days following the outpatient or emergency department visit.	Efficiency and Cost Reduction	Process High Priority	Orthopedic Surgery Physical Medicine General Practice/Family Medicine
eMeasure ID: CMS154v5 eMeasure NQF: N/A NQF: 0069 QualityID: 065	Appropriate Treatment for Children with Upper Respiratory Infection (URI)	Children age 3 months to 18 years who had an outpatient or emergency department (ED) visit with a diagnosis of upper respiratory infection (URI) during the measurement period.	Children without a prescription for antibiotic medication on or 3 days after the outpatient or ED visit for an upper respiratory infection.	Efficiency and Cost Reduction	Process High Priority	General Practice/Family Medicine Pediatrics
eMeasure ID: N/A eMeasure NQF: N/A NQF: N/A QualityID: 419	Overuse of Neuroimaging for Patients with Primary Headache and a Normal Neurological Exam	Patients with a diagnosis of primary headache.	Patients with a normal neurological examination for whom advanced brain imaging (CTA, CT, MRA, or MRI) was NOT ordered.	Efficiency and Cost Reduction	Efficiency High Priority	Neurology
eMeasure ID: N/A eMeasure NQF: N/A NQF: N/A QualityID: 333	Adult Sinusitis: Computerized Tomography (CT) for Acute Sinusitis (Overuse)	Patients aged 18 years and older with a diagnosis of acute bacterial sinusitis.	Patients who had a CT scan of the paranasal sinuses ordered at the time of diagnosis or received within 28 days after date of diagnosis.	Efficiency and Cost Reduction	Efficiency High Priority	Allergy/Immunology Internal Medicine Otolaryngology General Practice/Family Medicine

Compass PTN Core Measures

Measure ID	Measure Name	Denominator	Numerator	NQS Domain	Measure Type/ QPP Priority Level	QPP Specialty Measure Set
eMeasure ID: CMS50v5 eMeasure NQF: N/A NQF: N/A QualityID: 374	Closing the Referral Loop: Receipt of Specialist Report	Number of patients, regardless of age, who were referred by one provider to another provider, and who had a visit during the measurement period.	Number of patients with a referral, for which the referring provider received a report from the provider to whom the patient was referred.	Communication and Care Coordination	Process High Priority	Allergy/Immunology Cardiology Dermatology Emergency Medicine Gastroenterology General Surgery General Oncology Hospitalists Neurology Obstetrics/Gynecology Ophthalmology Orthopedic Surgery Otolaryngology Physical Medicine Preventive Medicine Rheumatology Thoracic Surgery Urology Vascular Surgery Mental/Behavioral Health Plastic Surgery
eMeasure ID: CMS68v6 eMeasure NQF: N/A NQF: 0419 QualityID: 130	Documentation of Current Medications in the Medical Record	All visits occurring during the 12 month reporting period for patients aged 18 years and older before the start of the measurement period.	Eligible professional attests to documenting, updating or reviewing the patient's current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosages, frequency and route of administration.	Patient Safety	Process High Priority	Allergy/Immunology Internal Medicine Anesthesiology Cardiology Dermatology Emergency Medicine Gastroenterology General Surgery General Oncology Hospitalists Neurology Obstetrics/Gynecology Ophthalmology Orthopedic Surgery Otolaryngology Physical Medicine Preventive Medicine Rheumatology Thoracic Surgery Urology Vascular Surgery Mental/Behavioral Health Plastic Surgery General Practice/Family Medicine

Compass PTN Core Measures

Measure ID	Measure Name	Denominator	Numerator	NQS Domain	Measure Type/ QPP Priority Level	QPP Specialty Measure Set
<p>NEW MEASURE eMeasure ID: N/A eMeasure NQF: N/A NQF: N/A QualityID: 402</p> <p><i>*This measure is a TCPI Common Measure</i></p>	Tobacco Use and Help with Quitting Among Adolescents	All patients aged 12-20 years with a visit during the measurement period	Patients who were screened for tobacco use at least once within 18 months (during the measurement period or the six months prior to the measurement period) AND who received tobacco cessation counseling intervention if identified as a tobacco user	Community/ Population Health	Process	Allergy/Immunology Internal Medicine Cardiology Dermatology Emergency Medicine Gastroenterology General Surgery General Oncology Hospitalists Neurology Obstetrics/Gynecology Ophthalmology Orthopedic Surgery Otolaryngology Physical Medicine Preventive Medicine Rheumatology Thoracic Surgery Urology Vascular Surgery Mental/Behavioral Health Plastic Surgery General Practice Family Medicine Pediatrics
<p>NEW MEASURE eMeasure ID: CMS2v6 eMeasure NQF: N/A NQF: 0418 QualityID: 134</p> <p><i>*This measure is a TCPI Common Measure</i></p>	Preventive Care and Screening: Screening for Clinical Depression and Follow-up Plan	All patients aged 12 years and older	Patients screened for depression on the date of the encounter using an age appropriate standardized tool AND , if positive, a follow-up plan is documented on the date of the positive screen	Community/ Population Health	Process	Internal Medicine Mental/Behavioral Health General Practice Family Medicine Pediatrics

Compass PTN Optional Measures

Measure ID	Measure Name	Denominator	Numerator	NQS Domain	Measure Type/ QPP Priority Level	QPP Specialty Measure Set
eMeasureID: N/A eMeasure NQF: N/A NQF: 1799 QualityID: 444	Medication Management for People with Asthma	Patients 5-64 years of age with persistent asthma and a visit during the measure period.	The number of patients who achieved a proportion of days (PDC) of at least 75% for their asthma controller medications during the measurement year.	Efficiency and Cost Reduction	Process High Priority	Allergy/Immunology General Practice/Family Medicine Pediatrics
eMeasure ID: CMS131v5 eMeasure NQF: N/A NQF: 0055 QualityID: 117	Diabetes: Eye Exam	Patients 18 - 75 years of age with diabetes with a visit during the measurement period	Patients with an eye screening for diabetic retinal disease. This includes diabetics who had one of the following: A retinal or dilated eye exam by an eye care professional in the measurement period or a negative retinal or dilated eye exam (no evidence of retinopathy) by an eye care professional in the year prior to the measurement period	Effective Clinical Care	Process	Internal Medicine Ophthalmology General Practice Family Medicine
eMeasure ID: CMS142v5 eMeasure NQF: N/A NQF: 0089 QualityID: 019	Diabetic Retinopathy: Communication with the Physician Managing Ongoing Diabetes Care	All patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed	Patients with documentation, at least once within 12 months, of the findings of the dilated macular or fundus exam via communication to the physician who manages the patient's diabetic care	Effective Clinical Care	Process High Priority	Ophthalmology
eMeasure ID: CMS167v5 eMeasure NQF: N/A NQF: 0088 QualityID: 018	Diabetic Retinopathy: Documentation of Presence or Absence of Macular Edema and Level of Severity of Retinopathy	All patients aged 18 years and older with a diagnosis of diabetic retinopathy	Patients who had a dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy AND the presence or absence of macular edema during one or more office visits within 12 months	Effective Clinical Care	Process	Ophthalmology

TCPI Common Measures

Measure ID	Measure Name	Denominator	Numerator	NQS Domain	Measure Type/ QPP Priority Level	QPP Specialty Measure Set
TCPI 01 eMeasure ID: N/A NQF: N/A QualityID: N/A	Documentation of a Comprehensive Health and Life Plan Developed Collaboratively by the Patient and the Health Professional Team	Total number of patients, of all ages, with two or more chronic conditions in each practice.	Number of patients, of all ages, with two or more chronic conditions in each practice for whom a comprehensive health and life plan is documented in the clinical record at each visit.	NQS Priority: Strengthen Person and Family Engagement	Process	
TCPI 02 eMeasure ID: N/A NQF: N/A QualityID: N/A	Referral of At-risk Patients to Community Based Prevention and Support Resources	Number of at-risk patients of all ages.	Number of at-risk patients, of all ages, referred to local community resources.	NQS Priority: Work with communities to promote best practices of healthy living	Process	
TCPI 03 eMeasure ID: N/A NQF: N/A QualityID: N/A	Medication Management	All patients regardless of age.	The number of patients, regardless of age, who received medication management services to include the following data: <ul style="list-style-type: none"> • Medications prescribed by any clinician • New medications to include: medications the patient has taken previously and are being re-started, medications the patient has not taken before • Any changes in medication regimen (change in dose, frequency, duration) • Any over the counter medications, vitamins, or herbal preparations • Review of allergies, including food-related allergies 	NQS Priority: Make Care Safer by Reducing Harm, Communication and Coordination of Care	Process	
eMeasure ID: CMS2v6 NQF: 0418 QualityID: 134 *This measure is a Compass PTN Core Metric	Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan	All patients aged 12 years and older.	Patient's screening for clinical depression using an age appropriate standardized tool AND follow-up plan is documented.	Community/ Population Health	Process	Internal Medicine Mental/Behavioral Health General Practice/Family Medicine Pediatrics
eMeasure ID: N/A NQF: N/A QualityID: 402 *This measure is a Compass PTN Core Metric	Tobacco Use and Help with Quitting Among Adolescents	All patients aged 12-20 years with a visit during the measurement period.	Patients who were screened for tobacco use at least once within 18 months (during the measurement period or the six months prior to the measurement period) AND who received tobacco cessation counseling intervention if identified as a tobacco user.	Community/ Population Health	Process	Allergy/Immunology Internal Medicine Cardiology Dermatology Emergency Medicine Gastroenterology General Surgery General Oncology Hospitalists Neurology Obstetrics/Gynecology Ophthalmology Orthopedic Surgery Otolaryngology Physical Medicine Preventive Medicine Rheumatology Thoracic Surgery Urology Vascular Surgery Mental/Behavioral Health Plastic Surgery General Practice/Family Medicine Pediatrics

TCPI Common Measures

Measure ID	Measure Name	Denominator	Numerator	NQS Domain	Measure Type/ QPP Priority Level	QPP Specialty Measure Set
eMeasure ID: N/A NQF: 2597 QualityID: N/A <i>*Measures in Composite NQF:0028; QualityID: 226 and NQF:2152; QualityID:431</i>	Substance Use Screening and Intervention Composite	All patients aged 18 years and older who were seen twice for any visits or who had at least one preventive care visit during the 12-month measurement period.	Patients who received the following substance use screenings at least once within the last 24 months AND who received an intervention for all positive screening results: <ul style="list-style-type: none"> • Tobacco use component <ul style="list-style-type: none"> o Patients who were screened for tobacco use at least once within the last 24 months AND who received tobacco cessation intervention if identified as a tobacco user; • Unhealthy alcohol use component <ul style="list-style-type: none"> o Patients who were screened for unhealthy alcohol use using a systematic screening method at least once within the last 24 months AND who received brief counseling if identified as an unhealthy alcohol user; and • Drug use component (nonmedical prescription drug use and illicit drug use) <ul style="list-style-type: none"> o Patients who were screened for nonmedical prescription drug use and illicit drug use at least once within the last 24 months using a systematic screening method AND who received brief counseling if identified as a nonmedical prescription drug user or illicit drug user. 	Community/ Population Health	Process	Allergy/Immunology Internal Medicine Cardiology Dermatology Emergency Medicine Gastroenterology General Surgery General Oncology Hospitalists Neurology Obstetrics/Gynecology Ophthalmology Orthopedic Surgery Otolaryngology Physical Medicine Preventive Medicine Rheumatology Thoracic Surgery Urology Vascular Surgery Mental/Behavioral Health Plastic Surgery General Practice/Family Medicine
eMeasure ID: CMS165v5 NQF: 0018 QualityID: 236 <i>*This measure is a Compass PTN Core Metric</i>	Controlling High Blood Pressure	Patients 18-85 years of age by the end of the measurement year who had at least one outpatient encounter with a diagnosis of hypertension (HTN) during the first six months of the measurement year.	The number of patients in the denominator whose most recent BP is adequately controlled during the measurement year. For a patient's BP to be controlled, both the systolic and diastolic BP must be <140/90mmHg (adequate control). To determine if a patient's BP is adequately controlled, the representative BP must be identified.	Effective Clinical Care	Intermediate Outcome High Priority	Internal Medicine Cardiology Obstetrics/Gynecology Preventive Medicine Thoracic Surgery Vascular Surgery General Practice/Family Medicine
eMeasure ID: CMS166v6 NQF: 0052 QualityID: 312 <i>*This measure is a Compass PTN Core Metric</i>	Use of Imaging Studies for Low Back Pain	Patients 18-50 years of age with a diagnosis of low back pain during an outpatient or emergency department visit.	Patients without an imaging study conducted on the date of the outpatient or emergency department visit or in the 28 days following the outpatient or emergency department visit.	Efficiency and Cost Reduction	Process High Priority	Orthopedic Surgery Physical Medicine General Practice/Family Medicine